

READ ENTIRE LABEL BEFORE USE



LUCROS-F™

F1837 - 11046 - 01 - Batch# - V021511OR

DESCRIPTION: Lucros-F is a foliar-applied liquid nutrient product that is designed to provide nutrients that may be deficient in the soil or plant.

DIRECTIONS FOR USE: Tank mix compatibility is impacted by water quality which may vary by location. Lucros-F is compatible with most of our other products. Lucros-F may be mixed with other foliar nutrient products, fungicides, insecticides, or herbicides. When mixing Lucros-F with other materials always establish compatibility using the standard quart jar method prior to tank mixing. Lucros-F should be applied as a foliar spray. Spray in the late evening or early morning. DO NOT spray to the point of runoff. Use as fine of a spray mist as possible. Allow 3-4 hours after application before sprinkler irrigation to avoid washing product off.

GUARANTEED ANALYSIS

Boron (B)..... 2.0%

Derived from: Boric Acid

2.50 US Gal / 9.46 Liters
8.35 lbs per Gallon @ 68° F
1.00 kg per Liter @ 20° C
Net Wt. 20.87 lb / 9.46 kg

SHAKE WELL BEFORE USE DO NOT STORE BELOW 40°F (5°C)

WARNING: This product contains Boron. Over application or excessive use of Boron, or use on crops other than those listed, can result in Boron toxicity, causing foliar damage or serious plant injury.

CAUTION: Keep Out Of Reach Of Children. Harmful if swallowed. In case of contact with eyes, flush immediately with copious amounts of water for 15 minutes. If irritation persists, consult a physician. Avoid contamination of skin and clothing. Wash with soap and water.

WARRANTY AND CONDITIONS OF SALE: Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. **BEFORE OPENING**, see other conditions and limitations that apply which are available at: <http://www.FBSciences.com/ProductConditionsOfSale/>.

Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.htm>

CROP TYPES	RATE (PER ACRE)	DETAILS
Sugar Beets	1 pint	Apply at full canopy (6-8 weeks prior to harvest) with 5-10 gallons of spray solution. A second application may be made 10-14 days later.